

Treatment of Hereditary Angioedema

Last updated March 2026

Treatment of Hereditary Angioedema: Replacement therapy or immune modulating medicines pertaining to hereditary angioedema.

Generic Name	Brand Name	Use
Berotrastat	Orladeyo Orladeyo Website	2 years and above as prophylaxis for hereditary angioedema
C1 Esterase Inhibitor (recombinant)	Ruconest Ruconest Website	Acute angioedema in hereditary angioedema
C1 Inhibitor (human)	Berinert Berinert Website	Acute attacks of hereditary angioedema Intravenous medication to treat acute attacks of hereditary angioedema.
	Cinryze Cinryze Website	To prevent angioedema due to C1 esterase inhibitor deficiency. Intravenous medication to prevent angioedema due to C1 esterase inhibitor deficiency.
	Haegarda Haegarda Website	Subcutaneous medication to prevent angioedema due to C1 esterase inhibitor deficiency.
Donidalorsen	Dawnzera Dawnzera Website	Indicated for prophylaxis to prevent attacks of hereditary angioedema (HAE) in adult and pediatric patients 12 years of age and older. The dose is 80 mg administered subcutaneously every 4 weeks.
Ecallantide Boxed Warning*	Kalbitor Kalbitor Website	Acute treatment of angioedema. From C1 esterase inhibitor deficiency. Three injections.
Garadacimab	Andembry Andembry Website	For prophylaxis of hereditary angioedema (type 1 and type 2). 12 years and above 400 mg loading dose followed by 200 mg subcutaneous every month.
Icatibant	Firazyr Firazyr Website	Acute attacks of hereditary angioedema- can be self-administered

		1 dose with onset of attack, can repeat in 6 hours, no more than 3 doses in 24 hrs
Lanadelumab	Takhzyro Takhzyro Website	For prophylaxis of hereditary angioedema (type 1 and type 2). 2-5 years 150 subcutaneous every 4 weeks. 6-11 years 150 mg subcutaneous every 2 weeks; consider every 4 weeks after 6 months. 12 years and above 300 mg subcutaneous every 2 weeks; consider every 4 weeks after 6 months.
Sebetralstat	Ekterly Ekterly Website	An oral on-demand therapy to treat hereditary angioedema (HAE) attacks in adults and children 12 years of age and older.

**Boxed Warning – this label indicates that the product, per the FDA, may have serious safety risks. Please speak with your health care provider or pharmacist to learn more about the potential side effects.*

© 2026 American Academy of Allergy, Asthma & Immunology. All Rights Reserved.